

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	▼ for					Go	Clear
Limits		Preview/Index		History		Clipboard	
Display		Summary		▼ Save		Text Order Add to Clipboard	
Show: 20		▼		Items 1-20 of 115		Page 1 of 6 Select page: 1 2 3 4 5 6	

Entrez PubMed

☐ 1: [Hatta H, Kim M, Yamamoto T.](#)

Related Articles

A novel isolation method for hen egg yolk antibody, "IgY".

Agric Biol Chem. 1990 Oct;54(10):2531-5.

PMID: 1368596 [PubMed - indexed for MEDLINE]

NEW

PubMed  
Services☐ 2: [Chang HM, Lu TC, Chen CC, Tu YY, Hwang JY.](#)

Related Articles

Isolation of immunoglobulin from egg yolk by anionic polysaccharides.

J Agric Food Chem. 2000 Apr;48(4):995-9.

PMID: 10775340 [PubMed - indexed for MEDLINE]

☐ 3: [Overfield ND.](#)

Related Articles

Eggstraordinary?

Vet Rec. 1995 Dec 23-30;137(26):676. No abstract available.

PMID: 8966981 [PubMed - indexed for MEDLINE]

Related  
Resources☐ 4: [Jensenius JC, Koch C.](#)

Related Articles

On the purification of IgG from egg yolk.

J Immunol Methods. 1993 Aug 26;164(1):141-2. No abstract available.

PMID: 8360505 [PubMed - indexed for MEDLINE]

☐ 5: [Gornall DA, Delahunty TJ, Kuksis A.](#)

Related Articles

Immunochemical relationships among plasma and egg-yolk lipoproteins of the laying hen.

Biochem J. 1971 Dec;125(4):97P-98P. No abstract available.

PMID: 4111387 [PubMed - indexed for MEDLINE]

☐ 6: [Suarez D.](#)

Related Articles

Incorporation of lycopene in egg yolk.

Poult Sci. 1969 Mar;48(2):733-5. No abstract available.

PMID: 5389891 [PubMed - indexed for MEDLINE]

☐ 7: [Mainguy P, Rouques A.](#)

Related Articles

[The yellow of the egg. I. General study of the color of the egg yolk].

Aliment Vie. 1965;53(4):83-116. Review. French. No abstract available.

PMID: 5322243 [PubMed - indexed for MEDLINE]

☐ 8: [Lin YC.](#)

Related Articles

A comparison of the nutritive value in three kinds of egg yolk.

Taiwan Yi Xue Hui Za Zhi. 1965 Nov 28;64(11):690-6. No abstract available.

PMID: 5218074 [PubMed - indexed for MEDLINE]

☐ **9:** MacKenzie SL, Martin WG.

Related Articles

The macromolecular composition of hen's egg yolk at successive stages of maturation.

Can J Biochem. 1967 Apr;45(4):591-601. No abstract available.

PMID: 6068014 [PubMed - indexed for MEDLINE]

☐ **10:** Hassl A, Aspöck H.

Related Articles

Purification of egg yolk immunoglobulins. A two-step procedure using hydrophobic interaction chromatography and gel filtration.

J Immunol Methods. 1988 Jun 13;110(2):225-8.

PMID: 3379312 [PubMed - indexed for MEDLINE]

☐ **11:** Ramachandran N, Kumar KS.

Related Articles

Characterization of phosvitin from hen egg yolk.

Indian J Biochem. 1967 Dec;4(4):210-1. No abstract available.

PMID: 4233073 [PubMed - indexed for MEDLINE]

☐ **12:** Bade H, Stegemann H.

Related Articles

Rapid method of extraction of antibodies from hen egg yolk.

J Immunol Methods. 1984 Sep 4;72(2):421-6.

PMID: 6432912 [PubMed - indexed for MEDLINE]

☐ **13:** James LK, Sherman P.

Related Articles

Creep compliance studies of egg yolk lipoprotein interfacial films.

Biorheology. 1976 Feb;13(1):79-81. No abstract available.

PMID: 945754 [PubMed - indexed for MEDLINE]

☐ **14:** Jones D.

Related Articles

Variations in the cholesterol content of egg yolk.

Nature. 1969 Feb 22;221(182):780. No abstract available.

PMID: 5766658 [PubMed - indexed for MEDLINE]

☐ **15:** Fischer M, Hlinak A, Montag T, Claros M, Schade R, Ebner D.

Related Articles

[Comparison of standard methods for the preparation of egg yolk antibodies].

Tierarztl Prax. 1996 Aug;24(4):411-8. German.

PMID: 9012029 [PubMed - indexed for MEDLINE]

☐ **16:** Steer DC, Martin WG, Cook WH.

Related Articles

Structural investigation of the low-density lipoprotein of hen's egg yolk using proteolysis.

Biochemistry. 1968 Sep;7(9):3309-15. No abstract available.

PMID: 5693359 [PubMed - indexed for MEDLINE]

☐ **17:** Dearden SJ, Hunter TF, Philp J.

Related Articles

A rapid method for the preparation of microvesicles of egg yolk lecithin.

Biochim Biophys Acta. 1982 Jul 28;689(2):415-8. No abstract available.

PMID: 6889439 [PubMed - indexed for MEDLINE]

☐ **18:** Burley RW.

Related Articles

A temperature-sensitive lipoprotein from abnormal egg yolk.  
Biochim Biophys Acta. 1970 Jul 14;210(2):347-9. No abstract available.  
PMID: 5476266 [PubMed - indexed for MEDLINE]

☐ 19: [Blount JD, Houston DC, Moller AP.](#)

[Related Articles](#)

Why egg yolk is yellow.  
Trends Ecol Evol. 2000 Feb;15(2):47-49. No abstract available.  
PMID: 10652553 [PubMed - as supplied by publisher]



☐ 20: [Vadehra DV, Bain JM, Burley RW.](#)

[Related Articles](#)

Lipid-protein globules of avian egg yolk. Isolation and properties of globules stable in concentrated sodium chloride solution.  
Biochem J. 1977 Sep 15;166(3):619-24.  
PMID: 563717 [PubMed - indexed for MEDLINE]

Display	Summary	▼	Save	Text	Order	Add to Clipboard
Show:	20	▼	Items 1-20 of 115		Page 1 of 6	Select page: 1 2 3 4 5 6

[Write to the Help Desk](#)  
[NCBI](#) | [NLM](#) | [NIH](#)  
[Department of Health & Human Services](#)  
[Freedom of Information Act](#) | [Disclaimer](#)



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	Clipboard
Search	PubMed	▼	for		Go	Clear	
Limits    Preview/Index    History    Clipboard							

Entrez PubMed

☐ 1: Dev Comp Immunol 1984 Fall;8(4):845-54[Related Articles, Books](#)

NEW

**Immunoglobulin classes in the garden lizard, *Calotes versicolor*.**PubMed  
Services**Natarajan K, Muthukkaruppan VR.**Related  
Resources

Two classes of immunoglobulins have been purified from lizard serum using a combination of ion-exchange chromatography on DEAE-cellulose and gel filtration on Sephadex G-200 or on Sepharose 6B. Lizard IgM is 2-ME sensitive and has an intact molecular weight similar to human IgM. On SDS-PAGE, reduced IgM dissociates into heavy and light chains of molecular weight 70,000 and 23,500 daltons respectively. Lizards also possess a 2-ME resistant, low molecular weight immunoglobulin designated as IgY similar to avian and amphibian IgY. IgY dissociates on SDS-PAGE to yield 59,500 dalton heavy and 26,000 dalton light chains. Antisera raised in rabbits to each of the two Ig classes could be made class-specific by cross-absorption, thus indicating that IgM and IgY represent distinct isotypes.

PMID: 6335107 [PubMed - indexed for MEDLINE]

Display	Abstract	▼	Save	Text	Order	Add to Clipboard
---------	----------	---	------	------	-------	------------------

[Write to the Help Desk](#)[NCBI](#) | [NLM](#) | [NIH](#)[Department of Health & Human Services](#)[Freedom of Information Act](#) | [Disclaimer](#)